



## Annual Research & Review in Biology

30(5): 1-9, 2018; Article no.ARRB.47022  
ISSN: 2347-565X, NLM ID: 101632869

# Physicochemical Characteristics of Snail as Affected by Processing Methods, Temperatures and Storage Days

I. Iwanegbe<sup>1\*</sup>, G. U. Emelue<sup>2</sup>, and J. U. Obaroakpo<sup>1</sup>

<sup>1</sup>Department of Food Technology, Auchi Polytechnic, Auchi, Edo State, Nigeria.

<sup>2</sup>Department of Forestry and Wildlife, Faculty of Agriculture, University of Benin, PMB 1154, Benin City, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. Author II designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors GUE and JUO managed the analyses of the study. Author JUO managed the literature searches. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/ARRB/2018/v30i530023

#### Editor(s):

- (1) Dr. Jean-Marie Exbrayat, Professor, Universite Catholique de Lyon, France.  
(2) Dr. George Perry, Dean and Professor of Biology, University of Texas at San Antonio, USA.

#### Reviewers:

- (1) Dr. O. A. Adedokun, Federal University of Technology, Nigeria.  
(2) Adeyeye, Samuel Ayofemi Olalekan, Ton Duc Thang University, Ho Chi Minh City, Vietnam.  
Complete Peer review History: <http://www.sdiarticle3.com/review-history/47022>

Original Research Article

Received 03 December 2018  
Accepted 12 February 2019  
Published 13 March 2019

## ABSTRACT

This study investigated the effect of processing methods, temperature and storage days on the physico-chemical characteristics of snail meat products. Four different treatments were carried out; unseasoned fried (USF), seasoned fried (SF), seasoned oven-dried (SOD) and seasoned smoke-dried (SSD) snail meat products and were stored under room, fridge and freezer storage conditions; physico-chemical determination and analysis of variance were carried out. The results showed that the highest crude protein was obtained in the seasoned smoke-dried product (76.87%), followed by seasoned oven-dried product (75.80%), next was seasoned fried (70.15%) and the least was unseasoned fried product (68.57%). The highest ash value was seen in seasoned smoke-dried product (4.84%), followed by seasoned oven-dried (4.313%) and the least was unseasoned-fried product (3.933%). The highest energy values were observed in the fried products (1497.67 KJ/100 g seasoned fried and 1490.53 KJ/100 g unseasoned-fried). The highest iron value (16.47 mg/100 g)

\*Corresponding author: E-mail: wanegbe@yahoo.com;

obtained was from seasoned smoke-dried product. Seasoned smoke-dried product was significantly different from other products and it had the highest mineral content. Seasoned oven-dried snail meat product had the lowest pH values 6.0, 7.17 and 6.29 for 0-5 days, 10-20 days and 25-30 days storage respectively. This was followed by seasoned smoke-dried product having 6.68, 7.81 and 6.56 at 0-5 days, 10-20 days and 25-30 days respectively. The observed low pH in seasoned products could be ascribed to the effect of the seasonings; this is an indication of better shelf stability. It was observed that the combine effect of seasonings, smoke-drying and cold storage help to extend the shelf life of snail meat.

**Keywords:** Processing; temperature; snail; products; seasoned.

## 1. INTRODUCTION

The land snails, particularly the *Archachatina marginata*, are alternative and non-conventional animal protein source in Nigeria and some other parts of Africa. Snails are processed by different methods for consumption. The different processing methods include roasting, oven-drying, frying etc. However, Onyeike and Oguike [1] and Ojiako, et al. [2] reported that different processing methods including boiling and roasting, influence the proximate, mineral and toxicant composition of foods. Preservation extends the shelf life of meat sample. Also, it is regarded as any method of treatment of food material to prolong the length of time in which it retains its qualities and appearance. Preservation methods include freezing, thermal processing (dehydration e.g. oven drying and smoke-drying) and curing method by using chemical additives [3].

Smoke-drying of meat is a technique whereby meat is exposed directly to wood smoke which may be generated by a variety of methods. There are various substances in the smoke produced from wood which contribute to the flavour and the appearance of the smoke-dried meat product and this has certain preserving effect on the product [3]. The preserving effect of smoke is not very significant when storing the product without a cold chain. On the other hand, intensive or prolonged smoke-drying may considerably increase the shelf-life of the product, but it also has an unfavourable effect on flavour. Whereas a light smoke generally enhances the organoleptic properties of the product, intensive smoking has a negative effect on the quality, especially in the case of prolonged storage in which concentrated smoke compounds develop increasingly unpleasant flavour.

In modern meat processing industries, beef, pork, lamb, veal and calf carcasses are chilled in chill cooler at temperature ranging from -4 to 0°C. Chilling storage is generally regarded to be

storage at temperature not far above freezing. The refrigerated storage of meat and meat products at 5°C-10°C is generally limited to relatively short periods, since deteriorative changes continue to occur. The major factors that influence the storage life of meat under refrigeration include the initial microbial load, temperature and humidity conditions during storage, the presence or absence of protective coverings, the species of animal involved, and the type of product being stored. Freezing as a preservation method is not a new process. It has long been recognized as an excellent means of meat preservation.

Snails are considered a delicacy in many countries and are staple part of the diet in parts of Asia where red meat and poultry are scarce sources of protein. In Nigeria, it is now accepted that the use of mini-livestock such as snails, rodents and other small livestock in the wild can substantially improve the living conditions of people in urban and rural areas by acting as a valuable source of protein supplement to diet as well as generating additional income [4].

The aim of this study was to investigate the effect of processing methods, temperature and storage days on the physico-chemical characteristics of snail meat products.

## 2. MATERIALS AND METHODS

### 2.1 Source of Snails

The snails used for this experiment were collected from Ekiuwa market in Edo State, Nigeria. A total of 150 adult snails (*Archachatina marginata*) with mean live weight of 346.85 g, were used. They were transferred to University of Benin where they were processed.

### 2.2 Removal of Meat from Shells

Snails were fasted for 24 hours in order to empty their guts and to reduce contamination during processing. Then the snails were weighed and

separated into meat, shell, waste and fluid. The meats were washed with alum to remove the slime and cut to uniform weight range of 50-55 g.

### **2.3 Application of Pickle in the Preservation of Fresh Snail Meat**

Meat may be preserved by dry curing or with a pickling solution. The ingredients used in curing and pickling are sodium nitrate, sodium nitrite, sodium chloride, sugar, citric acid or vinegar etc. Various methods are used: the meat may be mixed with dry ingredients; it may be soaked in pickling solution; pickling solution may be pumped or injected into the flesh; or a combination of these methods may be used.

In this study, snails were cured in a prepared pickle solution containing 1.5% salt, 1.5% sugar, 0.5% thyme, 0.30% nutmeg, 0.30% ginger, 1.50% red pepper, 0.05% sodium sorbate, 0.05% sodium tripolyphosphate, 0.50% curry, 1.50% onion (Table 1) for 24hours in refrigeration temperature, before processing snails (frying, smoke-drying and oven-drying). However, the control was devoid of spices before frying.

### **2.4 Processing Methods**

#### **2.4.1 Smoke-drying**

Pickle cured snail meat were skewered and smoke-dried at 80°C for 2 hours 15 minutes in a smoking kiln at Kilishi factory, Ekenwan Campus, Benin City, Nigeria. Each snail meat was spread out with stick in a traditional bush meat processing manner to increase the surface area of the meat exposed to smoke and heat. The meat samples were spread on racks in the smoking kiln to ensure uniform drying of the individual product. Initial weights of snail meat prior to smoke-drying were taken and weights after smoking were equally recorded.

#### **2.4.2 Frying**

Pickle cured snails were fried at 170°C for 30 minutes in a deep pan fryer with Soya oil (cholesterol free). 15 minutes into frying, meats were removed from oil, allowed to cool and weighed.

#### **2.4.3 Oven-drying**

Pickle cured snail meats were oven-dried at 90°C for 4 hours 30 min using table electric oven. The racks inside the oven were wrapped with foil paper before the meats were spread on

them. At every 45 minutes interval, meats were removed, allowed to cool and weighed.

### **2.5 Packaging**

Snail meat products were allowed to cool before packaging all the products including seasoned smoke-dried product that was skewered. All products were sealed in low density cellophanes with the use of sealing machine.

### **2.6 Storage Temperatures**

In this experiment, three storage temperatures were used.

- Room temperature (28.5°C)
- Refrigeration temperature (9.5°C)
- Freezer temperature (-12.5°C)

### **2.7 Storage Period**

Snail meat products were stored for total duration of 30 days and meat samples were withdrawn for analyses as follows.

- 0 day (control)
- 5 days
- 10 days
- 15 days
- 20 days
- 25 days
- 30 days

### **2.8 Analytical Methods**

Moisture content was determined by drying an accurately weighed sample of minced samples in an oven at 105±2°C for 3 hours. The ash content was obtained by heating the sample for 3 hours at 550°C. Fat was extracted according to the acid hydrolysis method. The total nitrogen content was determined by the Kjeldahl method and was converted to crude protein content by multiplying by 6.25 [5].

pH was determined on dispersion of two-gram sample in 10 ml of distilled water while a pocket pH meter was used to take pH values.

### **2.9 Data Analysis**

Data generated were subjected to analysis of Variance (ANOVA) to test significant variations ( $P<0.05$ ) among mean values obtained. Duncan's multiple range test was applied to indicate where significant differences ( $P<0.05$ ) occurred using Genstat statistical package 2005, 8<sup>TH</sup> edition (Genstat Procedure Library Release PL16).

**Table 1. Pickle formulation**

<b>Ingredients</b>	<b>Percentage (%)</b>	<b>Weight (g)</b>
Sugar	1.50	45
Salt	1.50	45
Thyme	0.50	15
Nutmeg	0.30	9
Ginger	0.30	9
Red pepper	1.50	45
Sodium sorbate	0.05	1.5
Sodium tripolyphosphate	0.05	1.5
Curry	0.5	15
Onion	1.50	45
Water	91.85	2755.5
Total	100%	3000 g

### 3. RESULTS AND DISCUSSION

#### 3.1 Chemical Composition of Snail Meat Products

The result of this study showed that the crude protein, fat, moisture, ash, fiber and carbohydrate of raw snail, were 16.69%, 4.87%, 64.03%, 3.78%, 3.47% and 6.89% respectively. The value of crude protein (16.69%) obtained is similar to the result (16.82%) of [6]. The moisture content (64.03%) obtained was similar to the result (63.1%) of Malik *et al.* (2011) but different from the work of Okonkwo and Anyaene [6] who had 79.48%. The crude fiber (3.47%) obtained was not different from the result (3.45 %) reported by Omoyakhi and Osinowo [7].

Analysis of variance showed significant difference ( $P<0.001$ ) in the chemical composition of the products based on treatment applied (Table 2). There was significant difference ( $P<0.05$ ) in the crude protein content among the products. The highest crude protein was obtained in the seasoned smoke-dried product (76.87%), followed by oven-dried product (75.80%), next was seasoned fried (70.15%) and the least was unseasoned fried product (68.57%). The high protein value in seasoned smoke-dried product demonstrates that smoke component has preservative influence on crude protein due to reduction effect on pH by smoke components. Besides, wood smoke contains pyrolygenous acid which may have added preservative effect on smoke-dried meat. Akhter *et al.* [8] reported that protein value of smoked meat (77.92%) product was significantly ( $P<0.05$ ) higher than crude protein of meat obtained from other processing methods. This could be attributed to lowering effect of pH by smoke components.

In terms of moisture content the different processing methods drastically reduced the moisture content of the raw snail meat. Iwanegbe, *et al.* [9] reported that processing methods have the potential of removing moisture from samples. Chima and Akobundu [10] reported that moisture content of sample decreases significantly ( $P<0.05$ ) with processing. The seasoned oven-dried product with the lowest moisture content was significantly different ( $P<0.05$ ) from other products. Smoke-dried product was significantly different ( $P<0.05$ ) from other products in terms of ash content. The highest ash value was seen in seasoned smoke-dried product (4.84%), followed by seasoned oven-dried (4.313%) and the least was unseasoned-fried (3.933%) although, Adegbite, *et al.* [11] reported ash content of 4.23% for snail of 6-12 months old.

For fat content, seasoned-fried (4.920%) and unseasoned-fried (5.033%) snail products were significantly different ( $P<0.05$ ) from smoke-dried (2.513%) and oven-dried products (2.57%). The high values recorded in the fried products could be attributed to oil absorption by the meat.

There was significant difference ( $P<0.05$ ) in carbohydrate content among all the products. The highest carbohydrate value was seen in unseasoned-fried product (8.16%) and the lowest was smoke-dried product (1.86%).

There was significant difference ( $P<0.05$ ) among the products in terms of crude fiber. However, the unseasoned-fried product had the highest fiber (3.77%) content. The highest energy values were seen in the fried products (1497.67 KJ/100 g seasoned fried and 1490.53 KJ/100 g unseasoned-fried). Although, all the products energy values ranged from 1431.40 - 1497.67 KJ/100 g, these values are lower than the energy

(1726 – 1740 KJ/100 g) reported by Engman, et al. [12] but greater than the values reported by Oduro, et al. [13] who reported 390.92 - 435.97 KJ/100 g. The energy values obtained in this study shows that snail meat could provide appreciable amount of calories in diet.

The results of this work showed that raw snail has Calcium (124.32 mg/100 g), iron (2.27 mg/100 g), Phosphorus (21.97 mg /100 g), Magnesium (23.95 mg/100 g), Copper (1.08 mg/100 g) and potassium (26.7 mg /100 g) contents. The values obtained are not different from the result of Malik et al. (2011) who reported Ca (126 mg/100 g), Fe (2.29 mg/100 g), P (22.9 mg/100 g), Mg (25.1 mg/100 g) and Cu (1.03 mg/100g). Adeola, et al. [14] reported Calcium (187 mg/100 g) and potassium (25.6 mg/100 g) for raw snail. The minerals for processed snail range from 146.3 mg/100 g-165.7 mg/100 g for calcium, 14.07-16.47 mg/100 g iron, 183.0-235.0 mg/100 g phosphorus, 74.63-104.23 mg/100 g copper, 305-386.3 mg/100 g potassium and 57.63-64.63 mg/100 g magnesium.

From the analysis of variance the seasoned smoke-dried product was significantly different ( $P<0.05$ ) from other products and had the highest mineral content. The value obtained for P (235.0 mg/100 g) was not different from the value (238.0 mg/100 g) obtained by Kalio, et al. [15]. Phosphorus and potassium are important in human and animal nutrition. Phosphorus is used for normal development and maintenance of bones and teeth, cell activity, normal acid-base balance of blood, muscle activity, metabolism of carbohydrate and fat.

The highest iron value obtained (16.47 mg/100 g) was from seasoned smoke-dried product. This is close to the result (12.2 mg/100 g) obtained by [16]. However, the value obtained was not different from 16.1 mg/100g reported by Kalio, et al. [15]. Wosu [16] reported that iron content of snail varies from one locality to another depending on mineral content of the soil in which these snails are raised. Iron is good for bone and teeth formation as well as for haemoglobin of the red blood cells. Cobalt (Co) was not detected. According to Fagbua, et al. [17] the non-detection of lead and cobalt confirm that none of the snail had been exposed to any sort of pollution.

### 3.2 pH Changes of Snail Meat Products

The analysis of variance showed that there was significant difference ( $P<0.001$ ) in the main and

interactive effects in pH values of the different snail meat products based on treatments (Processing methods, storage conditions and storage days).

The average pH value of raw snail washed without alum was 7.02 while raw snail washed with alum had an ultimate pH of 7.3. This value agreed closely with the work of [6] who reported pH value of 7.4, this according to them is due to the basicity of potassium alum used for washing the foot, which tends to raised pH value.

Table 3 showed the main effect of processing methods on the pH of products. There was significant difference ( $P<0.05$ ) in the different products stored for 0-5 days, 10-20 days and 25-30 days. Seasoned oven-dried snail meat product (412) had the lowest pH values 6.0, 7.17 and 6.29 for 0-5 days, 10-20 days and 25-30 days storage respectively. This was followed by seasoned smoke-dried product having 6.68, 7.81 and 6.56 respectively. Moreover, it was observed that seasoned products had lower pH than the unseasoned product, an indication of better shelf stability of the seasoned products. This could be attributed to the presence of salt and other curing ingredients which altered the pH of the seasoned products, thereby limiting the growth of spoilage organisms. The result of this study showed that the interactive effect of salt and other spices inclusion in the seasoned snail meat products lowered the pH thereby ensuring shelf stable products.

Table 4 showed the effect of storage conditions on pH of products. There was significant difference ( $P<0.05$ ) between products under room storage (7.20) and products under cold storage (fridge 6.61 and freezer 6.05) at 5 days. Products under room and fridge storage could not last beyond 5 days and 20 days respectively. Table 5 also showed significant difference ( $P<0.05$ ) in the pH of products stored for 10 days (7.51), 15 days (8.22) and 20 days (8.59).

Table 6 showed the pH values of the interaction between processing methods and storage periods. The pH values of the snail meat products significantly ( $P<0.05$ ) increased with the storage days (0-5 and 10-20) but with insignificant ( $P>0.05$ ) decline at 25 days. Seasoned oven-dried product had the lowest pH values of 5.76, 6.69 and 6.26 for 0-5 days, 10-20 days and 25-30 days storage period respectively. This was followed by seasoned smoke-dried product (6.40, 7.54 and 6.49). The unseasoned-

fried product (control) had the highest pH values at 5 days (7.98), 20 days (9.77) and 25 days (8.33) storage. This high value was responsible for the short shelf life of the unseasoned fried product (control). Kiers, et al. [18] reported that the increase in pH value during storage is due to the degradation of protein.

**Table 2. Means for chemical composition of snail meat products**

Parameter	Unseasoned fried (311)	fried (312)	Seasoned oven-dried (412)	Seasoned smoke-dried (512)	LSD
Protein(%)	68.567 <sup>d</sup>	70.150 <sup>c</sup>	75.803 <sup>b</sup>	76.867 <sup>a</sup>	0.1770
Moisture (%)	10.543 <sup>a</sup>	10.050 <sup>b</sup>	9.563 <sup>c</sup>	10.467 <sup>a</sup>	0.0773
Ash (%)	3.933 <sup>d</sup>	3.963 <sup>c</sup>	4.313 <sup>b</sup>	4.837 <sup>a</sup>	0.0129
Fat (%)	5.033 <sup>a</sup>	4.920 <sup>a</sup>	2.570 <sup>b</sup>	2.513 <sup>b</sup>	0.1227
Crude fiber (%)	3.767 <sup>a</sup>	3.677 <sup>b</sup>	3.490 <sup>c</sup>	3.453 <sup>d</sup>	0.0179
Carbohydrate(%)	8.157 <sup>a</sup>	7.240 <sup>b</sup>	4.260 <sup>c</sup>	1.863 <sup>d</sup>	0.2681
Energy(KJ/100g)	1490.53 <sup>b</sup>	1497.67 <sup>a</sup>	1456.17 <sup>c</sup>	1431.40 <sup>d</sup>	2.813
Ca (mg/100g)	146.3 <sup>c</sup>	153.7 <sup>bc</sup>	156.7 <sup>b</sup>	165.7 <sup>a</sup>	8.22
Fe(mg/100g)	14.07 <sup>d</sup>	14.70 <sup>c</sup>	15.20 <sup>b</sup>	16.47 <sup>a</sup>	0.38
P (mg/100g)	183.0 <sup>d</sup>	196.3 <sup>c</sup>	204.7 <sup>b</sup>	235.0 <sup>a</sup>	7.33
Cu (mg/100g)	74.63 <sup>d</sup>	87.0 <sup>c</sup>	96.67 <sup>b</sup>	104.23 <sup>a</sup>	4.05
K (mg/100g)	305 <sup>d</sup>	327.3 <sup>c</sup>	358 <sup>b</sup>	386.3 <sup>a</sup>	10.98
Mg (mg/100)	57.63 <sup>c</sup>	60.55 <sup>b</sup>	62.97 <sup>a</sup>	64.63 <sup>a</sup>	2.29

Means with same superscript along the row are not significantly differently ( $P>0.05$ )

**Table 3. pH of snail meat products (processing methods)**

Storage days	Treatments (processing methods)				LSD
	Unseasoned/ fried	Seasoned/ fried	Seasoned/ oven-dried	Seasoned/ smoke-dried	
0 – 5	7.711 <sup>a</sup>	6.811 <sup>a</sup>	6.000 <sup>d</sup>	6.689 <sup>c</sup>	0.0839
10 – 20	9.147 <sup>a</sup>	8.046 <sup>b</sup>	7.170 <sup>d</sup>	7.817 <sup>c</sup>	0.1317
25 – 30	8.226 <sup>a</sup>	6.708 <sup>b</sup>	6.290 <sup>d</sup>	6.563 <sup>c</sup>	0.0207

Means within storage day bracket having same superscript are not significantly different ( $P>0.05$ )

**Table 4. pH of snail meat products (storage conditions)**

Storage days	Treatments(storage conditions)			LSD
	Room (28.5°C)	Fridge (9.5°C)	Freezer(-12.5°C)	
0 – 5	7.207 <sup>a</sup>	6.617 <sup>b</sup>	6.058 <sup>b</sup>	0.0727
10 – 20	-	9.129	6.961	.0927
25 – 30	-	-	-	-

Means within storage day bracket having same superscript are not significantly different ( $P>0.05$ )

**Table 5. pH of snail meat products (storage days)**

Storage days	Means values	LSD
0	6.525	0.0593
5	7.081	
10	7.518 <sup>c</sup>	
15	8.220 <sup>b</sup>	
20	8.595 <sup>a</sup>	
25	7.022	0.014
30	6.871	

Means with the same letters are not significantly different ( $P>0.05$ )

**Table 6. Effect of storage on pH changes of snail meat**

Products	Storage period (days)						
	0	5	10	15	20	25	30
311	7.433 <sup>b</sup>	7.989 <sup>a</sup>	8.278 <sup>d</sup>	9.385 <sup>b</sup>	9.778 <sup>a</sup>	8.333 <sup>a</sup>	8.120 <sup>b</sup>
312	6.500 <sup>e</sup>	7.122 <sup>c</sup>	7.557 <sup>gh</sup>	7.945 <sup>ef</sup>	8.635 <sup>c</sup>	6.813 <sup>c</sup>	6.603 <sup>d</sup>
412	5.767 <sup>g</sup>	6.233 <sup>f</sup>	6.697 <sup>j</sup>	6.977 <sup>i</sup>	7.837 <sup>f</sup>	6.313 <sup>f</sup>	6.266 <sup>g</sup>
512	6.400 <sup>e</sup>	6.978 <sup>d</sup>	7.540 <sup>h</sup>	7.782 <sup>ig</sup>	8.130 <sup>de</sup>	6.630 <sup>d</sup>	6.496 <sup>e</sup>
SEM	0.0417		0.0801		0.0096		

Means within storage day bracket having same superscript along the row and down the column are not significantly different ( $P>0.05$ ).

311= unseasoned fried, 312=seasoned fried, 412=seasoned oven-dried, 512=seasoned smoke-dried

**Table 7. pH of snail meat products (storage conditions and storage days)**

Storage Condition	Storage period (days)						
	0	5	10	15	20	25	30
Room (28.5°C)	6.525 <sup>c</sup>	7.883 <sup>a</sup>	-	-	-	-	-
Fridge (9.5°C)	6.525 <sup>c</sup>	6.708 <sup>b</sup>	8.285 <sup>c</sup>	9.168 <sup>b</sup>	9.932 <sup>a</sup>	-	-
Freezer (-12.5°C)	6.525 <sup>c</sup>	6.650 <sup>b</sup>	6.751 <sup>e</sup>	6.876 <sup>e</sup>	7.257 <sup>d</sup>	7.022	6.8717
SEM	0.0361		0.0567		0.0048		

Means within storage day bracket having same superscript along the row and down the column are not significantly different ( $P>0.05$ ).

**Table 8. pH of snail meat products (processing methods and storage conditions)**

Products	Fridge (9.5°C)	Freezer (-12.5°C)	Room (28.5°C)
<b>0-5 days</b>			
311	7.517 <sup>b</sup>	7.483 <sup>bc</sup>	8.133 <sup>a</sup>
312	6.533 <sup>e</sup>	6.550 <sup>e</sup>	7.350 <sup>c</sup>
412	5.817 <sup>h</sup>	5.817 <sup>h</sup>	6.183 <sup>f</sup>
512	6.417 <sup>e</sup>	6.500 <sup>e</sup>	7.150 <sup>d</sup>
SEM	0.0511	0.0511	0.511
<b>10-20 days</b>			
311	10.208 <sup>a</sup>	8.087 <sup>d</sup>	-
312	9.267 <sup>b</sup>	6.824 <sup>e</sup>	-
412	8.162 <sup>d</sup>	6.178 <sup>f</sup>	-
512	8.878 <sup>c</sup>	6.757 <sup>e</sup>	-
SEM	0.0654	0.0654	-
<b>25-30 days</b>			
311	-	8.227 <sup>a</sup>	-
312	-	6.708 <sup>b</sup>	-
412	-	6.290 <sup>d</sup>	-
512	-	6.563 <sup>c</sup>	-
SEM	0.0068	-	-

Means within storage day bracket having same superscript along the row and down the column are not significantly different ( $P>0.05$ ).

311= unseasoned fried, 312=seasoned fried, 412=seasoned oven-dried, 512=seasoned smoke-dried

Table 7 showed the changes in pH values of products due to the interaction between storage conditions and storage days. There was significant difference ( $P<0.05$ ) between pH of products stored for 0-5 days, 10-20 days and 25-30 days under the different storage conditions. Products under room storage increased significantly ( $P<0.05$ ) in pH than snail meat products under cold storage from 0-5 days. Also, products under fridge (6.70) storage condition

were not significantly ( $P>0.05$ ) different from snail meat under freezer (6.65) storage at 5 days. This implies that cold storage helps to control and stabilize pH of meat products thereby enhancing their shelf stability. Iwanegbe, et al. [9] reported that refrigeration extends shelf stability and prevent product deterioration. A significant increase ( $P<0.05$ ) was observed in the pH values of products under refrigerated storage from 10-20 days.

**Table 9. pH means of snail meat products (processing methods, storage days and storage conditions)**

Products storage conditions	Storage period(days)						
	0	5	10	15	20	25	30
311 Freezer(-12.5 <sup>o</sup> C)	7.433 <sup>d</sup>	7.533 <sup>d</sup>	7.703 <sup>gh</sup>	7.903 <sup>g</sup>	8.653 <sup>ef</sup>	8.333 <sup>a</sup>	8.120 <sup>d</sup>
312 Freezer(-12.5 <sup>o</sup> C)	6.500 <sup>e</sup>	6.600 <sup>e</sup>	6.670 <sup>i</sup>	6.867 <sup>i</sup>	6.937 <sup>j</sup>	6.813 <sup>c</sup>	6.603 <sup>d</sup>
412 Freezer(-12.5 <sup>o</sup> C)	5.767 <sup>g</sup>	5.867 <sup>g</sup>	5.933 <sup>j</sup>	6.027 <sup>j</sup>	6.313 <sup>f</sup>	6.573 <sup>i</sup>	6.266 <sup>g</sup>
512 Freezer(-12.5 <sup>o</sup> C)	6.400 <sup>ef</sup>	6.600 <sup>e</sup>	6.697 <sup>i</sup>	6.707 <sup>i</sup>	6.867 <sup>i</sup>	6.630 <sup>d</sup>	6.496 <sup>e</sup>
311Fridge(9.5 <sup>o</sup> C)	7.433 <sup>d</sup>	7.600 <sup>d</sup>	8.853 <sup>de</sup>	10.867 <sup>a</sup>	10.903 <sup>a</sup>	-	-
312 Fridge(9.5 <sup>o</sup> C)	6.500 <sup>e</sup>	6.567 <sup>e</sup>	8.443 <sup>f</sup>	9.023 <sup>d</sup>	10.333 <sup>b</sup>	-	-
412 Fridge(9.5 <sup>o</sup> C)	5.767 <sup>g</sup>	6.233 <sup>f</sup>	7.460 <sup>h</sup>	7.927 <sup>g</sup>	9.100 <sup>cd</sup>	-	-
512 Fridge(9.5 <sup>o</sup> C)	6.400 <sup>ef</sup>	6.433 <sup>ef</sup>	8.383 <sup>f</sup>	8.857 <sup>de</sup>	9.393 <sup>c</sup>	-	-
311 Room(28.5 <sup>o</sup> C)	7.433 <sup>d</sup>	8.833 <sup>a</sup>	-	-	-	-	-
312 Room(28.5 <sup>o</sup> C)	6.500 <sup>e</sup>	8.200 <sup>b</sup>	-	-	-	-	-
412 Room(28.5 <sup>o</sup> C)	5.767 <sup>g</sup>	6.600 <sup>e</sup>	-	-	-	-	-
512 Room(28.5 <sup>o</sup> C)	6.400 <sup>ef</sup>	7.900 <sup>c</sup>	-	-	-	-	-
SEM		0.0722			0.1133		0.0096

Means within storage day bracket having same superscript along the row and down the column are not significantly different ( $P>0.05$ ).

311= unseasoned fried, 312=seasoned fried, 412=seasoned oven-dried, 512=seasoned smoke-dried

The interaction between processing methods and storage conditions is shown in Table 8. At 0-5 days' storage, the pH of the various products at refrigeration condition was lower than products under room storage but higher than products under freezer storage. The snail meat product is better shelf stable under fridge and freezer storage due to lower pH as product in refrigeration storage had shelf life of 20 days, freezer storage could last 30 days and beyond while room stored products lasted for 6 days. This was also reflected in Table 9.

#### 4. CONCLUSION

The various processing methods caused reduction in the moisture content of products particularly oven-drying. The high protein value in seasoned smoke-dried product demonstrated that smoke component has preservative influence because of the polyphenols which has antimicrobial properties. All the seasoned products had low pH values than the unseasoned product and this could be ascribed to the effect of the seasonings. The pH of the product under refrigeration condition was lower than product under room storage but higher than product under freezer storage. For extended shelf life and increase in nutritive component of snail meat there should be a combined effect of seasonings, smoke-drying and cold storage.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Onyeike EN, Oguike JU. Influence of heat processing methods on the nutrient composition and lipid characterization of groundnut (*Arachis hypogaea*) seed pastes. *Biokemistri*. 2003;15(1):34-43.
2. Ojiako OA, Ogbuji CA, Agha NC, Onwuliri AVA. The proximate, mineral and toxicant composition of four possible food security crops from southern Nigeria. *J. Med. Food*. 2010;13(5):1203-1209.
3. Iwanegbe I, Igene JO, Emelue GU, Obaroakpo JU. Effect of processing, storage days and storage temperatures on lipid oxidation and palatability processed snail meat products. *Asian Food Science Journal*. 2019;6(2):1-12.
4. Ezeama CF, Keke E, Nwachukwu E. Influence of brime treatment, drying methods and storage conditions on the microbial quality of freshwater snail (*Lanistes libycus*) meat. *Nig Food J*. 2007; 25:101-111.
5. AOAC (Association of Official Analytical Chemists) Official Method of Analysis, 4<sup>th</sup> Ed; 2000.
6. Okonkwo TM, Anyaene LU. Meat Yield and the effects of curing on the characteristics of snail meat. *J. of Tropical Agriculture, Food, Environment and Extension*. 2009; 8(1):66-73.
7. Omoyakhi JM, Osinowo OA. Modification of some biochemical activities response to

- transition of giant African land snails. *Archachatina marginata* and *Achantina achantina* from aestivation to an Active state. Archives of Applied Science Research. 2010;2(3):53-60.
8. Akhter S, Rahaam M, Hossain MM, Hashern MA. Effect of drying as Preservation techniques on nutrient content of beef. J. Bangladesh Agril. Univ. 2009;7(1):163 –168.
  9. Iwanegbe I, Obaroakpo JU, Suleiman M. Effect of processing and storage methods on water content and activity of snail meat products. The Nigerian Journal of Agric. And Forrestry. 2018;6(1):59-70.
  10. Chima JU, Akobundu ENT. Proximate composition of processed freshwater snail (*Pilaovata*) meat as affected by salting, fermentation and frying. Journal Agric Environ. 2010;12(2);150 -156.
  11. Adegbite JA, Sanni LO, Osinowo OA. Comparative evaluation of chemical and sensory properties of *Achatina achatina* and *Archachatina marginata*. Assert an international Journal. Asset Series A. 2006;6(2):1-6.
  12. Engman FN, Ellis WO, Dzugbefia VP, Yong – Kim M, Abano E, Owusu J. A comparative study of three drying methods for preservation of the giant African snail (*Achatina achatina*) meat. African Journal of Food Science. 2012;6(14):392-400.
  13. Oduro W, Ellis WO, Oduro I, Tetteh D. Meat yield and quality of selected snail species in Ghana, Journal of Ghana Science Association. 2002;4(2):24-30.
  14. Adeola AJ, Adeyemo AI, Ogunjobi JA, Alaye SA, Adelakun KM. Effect of natural and concentrate diets on proximate composition and sensory properties of Giant Land Snail (*Archachatina marginata*) Meat. Journal of Applied Science in Environment Sanitation. 2010;5(2):185-189.
  15. Kalio GA, Godfrey AK, Ibisime E. Nutritional and sensory profiling of the African Giant land snail fed commercial type and leaf based diets in a rain-forest ecology. Journal of Food, Agric. Nutri. and Development. 2011;11(5).
  16. Wosu LO. Commercial snail farming in West African – A Guide Ap Express Publishers, Nsukka – Nigeria; 2003.
  17. Fagbuaro O, Oso JA, Edward JB, Ogunleye RF. Nutritional status of four species of giantland snails. Nigeria J. Zhejiang Univ. Sci. 2006;7(9):686– 689.
  18. Kiers JL, Van-Laeken AEA, Rombout FM, and Nout MJR. *In vitro* digestibility of Bacillus fermented soya bean. Int. J. Food Microbial. 2000;60:163 -169.

© 2018 Iwanegbe et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/47022>