

## Annual Research & Review in Biology

30(5): 1-16, 2018; Article no.ARRB.46533  
ISSN: 2347-565X, NLM ID: 101632869

# Phylogenetic Framework and Metabolic Genes Expression Analysis of Bacteria Isolated from Contaminated Marine Environments of Niger Delta

Bright Obidinma Uba<sup>1\*</sup>

<sup>1</sup>Department of Microbiology, Chukwuemeka Odumegwu Ojukwu University, P.M.B. 02 Uli, Anambra State, Nigeria.

### Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

### Article Information

DOI: 10.9734/ARRB/2018/v30i530023

#### Editor(s):

(1) Dr. George Perry, Dean and Professor of Biology, University of Texas at San Antonio, USA.

#### Reviewers:

(1) Noor Ul Akbar, Kohat University of Science and Technology, Pakistan.

(2) Dr. M. Thangaraj, Annamalai University, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46533>

Received 01 November 2018

Accepted 17 January 2019

Published 13 March 2019

Original Research Article

## ABSTRACT

**Aims:** To explore the phylogenetic framework of bacteria isolated from contaminated marine environments of Niger Delta and the expression of the metabolic genes coding for aromatic hydrocarbon degradation and surfactant production.

**Study Design:** Nine treatments designs were set up in triplicates containing 25 mL of sterile modified mineral basal medium supplemented with nine marine hydrocarbon degraders incubated at 24°C for 5 days. Three of the set ups were supplemented each with 1 mg /L of xylene, anthracene and pyrene.

**Place and Duration of Study:** Department of Environmental Sciences, University of South Africa, Pretoria, South Africa between September, 2015 to December, 2017.

**Methodology:** A laboratory scale study was carried on six composite samples of the sediment and water samples from the three studied areas using enrichment, screening, selection, characterization, and PCR assays to explore the phylogenetic framework and metabolic genes expression of the marine bacteria for aromatic hydrocarbon degradation and surfactant production.

**Results:** The findings revealed that there was significant abundance of THB ( $P = .05$ ) more than TCHUB and more xylene degraders than anthracene and pyrene degraders in the sediment and

\*Corresponding author: E-mail: [ubabright4real@yahoo.com](mailto:ubabright4real@yahoo.com);

water samples respectively. The phylogenetic correlational analysis revealed that all the nine selected best degraders out of 48 isolates from the studied area were evolutionary related belonging to the genera: *Providencia*, *Alcaligenes*, *Brevundimonas*, *Myroides*, *Serratia*, and *Bacillus*; able to significantly ( $P = .05$ ) utilize the all the aromatic hydrocarbons. The existence of catabolic and surfactant genes namely catechol dioxygenase (C23O), rhamnolipid enzyme (*rhlB*) and surfactin /lichenysin enzyme (*SrfA3 /LicA3*) genes were detected in only four (4) out of the nine (9) marine aromatic degrading bacteria with 881 base pairs sizes.

**Conclusion:** Thus, the study revealed that these bacterial strains especially *Serratia marcescens* XYL7 might possess metabolic genes for *in situ* aromatic hydrocarbon degradation and surfactant production.

**Keywords:** Aromatic hydrocarbons; biodegradation; biodiversity; catabolic genes; marine bacteria; Niger Delta; surfactant genes.

## 1. INTRODUCTION

The Niger Delta region is the centre of petroleum exploration, production and development accomplishments in Nigeria. Several cases of oil spills have been reported thereby promoting and increasing the pollution of the Niger Delta's aquatic and terrestrial environments. These spills have been associated with disruption, deterioration of pipes, negligence, mishaps during oil exploitation, production, rupture of storage tanks and accidents brought about by transportation [1]. Most of these compounds especially the aromatic hydrocarbons are considered as toxicants and are classified as priority environmental pollutants by the US Environmental Protection Agency [2].

Biodiversity is the result of evolution that occur billions of years ago leading to multiplicity of species of living organisms on the earth. Molecular phylogenetic studies have shown that the chief diversity of life is in the microbial world and it is circulated amongst the three main domains: eubacteria, eukarya, and archaea. The methods of single cell sequencing, metatranscriptomics, metaproteomics and metagenomics are used not only to reveal the community structure (species taxonomy, abundance and distribution) but also of the useful and biological niche of a community [3]. Most of biotechnological products such as biosurfactants, antibiotics, compatible solutes, exopolysaccharides, enzymes etcetera with enormous medical, environmental and industrial uses are obtained from microbes of marine ecosystems or sources. This has attracted significant attention and interest on the marine bacteria as they produces novel metabolites of diverse applications than the terrestrial and other sources. Several reports have implicated marine bacteria in the production of metabolites such as

surfactant that have purposeful and useful roles in bioremediation especially in the aspect of enhanced oil recovery [4].

As polluted sites are contaminated by a mixture of PAHs and other environmental contaminants [4], considerable investigation has concentrated on the bioremediation of aromatics hydrocarbons (Ahs) through metabolism and co-metabolism. The mechanism of degradation process chiefly involves enzyme machinery: dehalogenases, dehydrogenases, oxygenases and hydrolases system [5]. Thus, for an effectual remediation process to exist, it is imperative that the bacteria involved should have a comprehensive degradation pathway so that no possible noxious metabolites accrue [6]. Researches have shown that a cluster of genes for whole degradation of aromatic compounds are found in several PAH - degrading bacteria through genetic analyses of PAH catabolic pathways. The diversity of aromatic-dioxygenase genes in several PAH degrading bacterial isolates have been studied through PCR amplification using genes specific primers or degenerate primers to revealed them [4]. Even though numerous bacteria capable of degrading PAHs have already been isolated and documented, it is still important to search, monitor and select more efficient and potent strains especially of the marine origin that can absolutely and rapidly mineralize PAHs under the deplorable and unfriendly environmental conditions of Niger Delta. Thus, exploring the phylogenetic markers such as 16S rRNA genes to disclose the microbial diversity and further investigation of gene expression to unveil the metabolic power of the marine microbes in the Niger Delta is essential. Therefore, the aim of the present study was to explore the phylogenetic framework of bacteria isolated from contaminated marine environments of Niger Delta and the expression of the metabolic genes

coding for aromatic hydrocarbon degradation and surfactant production.

## 2. MATERIALS AND METHODS

### 2.1 Description of the Sampling Sites

The studied areas were Abonema Wharf Water Front (Fig. 1) in Akuku-Toru Local Government Area, Nembe Water-side (Fig. 2) in Port Harcourt Local Government Area and Onne Light Flow Terminal Seaport (Fig. 3) located in Eleme Local Government Area of Rivers State. Abonema town is 53 Km and Abonema Wharf Water Front is 3 - 5 Km from Port Harcourt capital city; Nembe water side is located within Port Harcourt capital city of Rivers State, while Onne Light Flow Terminal is about 35 Km east from Port Harcourt capital city of Rivers State and 7 km from Onne town. These sites were geo - referenced using Handheld Global Positioning System (GPS) GPSMAP 76 sc with the coordinates obtained from the sampling points or positions. Abonema Wharf Water Front, Nembe Water-side and Onne Light Flow Terminal Seaport were located between latitude 4°46'15.82"N to latitude 4°46'38.01"N and longitude 7°0'0.54"E to longitude 7°0'34.82"E with average elevation of 4.1 m , latitude 4°45'8.72"N to latitude 4°45'26.42"N and longitude 7°1'11.37"E to longitude 7° 2'14.54"E with average elevation of 2.7 m and latitude 4°41'32.58"N and 4°41'58.18"N and longitude 7°9'26.34"E and 7°10'48.82"E with average elevation of 2.3 m, respectively. These water - ways are subjected to anthropological - induced stresses brought about by expansion, industrial development and rigorous transportation. Abonema Wharf Water Front community is a popular and busy commercial but dangerous jetty area close to Port Harcourt city inhabiting tens of thousands of different families living close to petroleum tank farms and tankers queue up daily to load refined petroleum products. Nembe Waterside is located adjacent to Creek Road Market, Port Harcourt, Nigeria. It shares boundary with Bayelsa and connects Port Harcourt city with Bonny Island where greater percentage of petroleum connections in Rivers State are located. Also, it connects the Island straight to the Atlantic Ocean where petroleum is shipped by gigantic oil vessels [7]. Onne Light Flow Terminal Seaport is a port of Nigeria and the leading petroleum and gas unrestricted zone in the world supportive of exploration and production for Nigerian accomplishments. It is located on the Bonny River Estuary beside Ogu

Creek and accounts for more than 65% of the export cargo through the Nigerian Sea Port. Anthropological survey revealed the presence of human activities such as transportation of petrochemical products through tankers, canoes, boats and ships to neighboring villages, towns, cities, states and nations due to the presence of multinational petrochemical and oil servicing industries such as Chevron Nigeria Limited, Cameron Offshore services, Exxon Mobil Nigeria Limited, Socotherm Pipecoaters, Beker Hughes Oil Servicing Company, Aiteo Energy Resource, Sorelink Oil and Dozzy Oil and gas etcetera that generate the wastes that contaminate the sites above.

### 2.2 Collection and Processing of Sample

Ten samples each of the marine sediment and water were collected randomly per point of the designated ten (10) points of the three sampling sites (Figs. 1, 2 and 3). The samplings were done once in each of the three sampling sites in September, 2015. The samples were mixed together after which a total of six composite /representative sediment and water samples were obtained and taken for the analysis. The surface aerobic sediment samples were collected with a 95% ethanol - sanitized plastic spatula at 5 cm depth inside 95% ethanol - sanitized wide mouthed plastic containers. The water samples were collected at the air-water interface by hand dipping the 95% ethanol - sanitized cylindrical shaped 2 L plastic containers. The containers were rinsed with the sediment and water samples before collecting the samples. All the composite or representative sediment and water samples containers were placed into a sterile polythene bag and then transported to the Microbiology Laboratory, Chukwuemeka Odumegwu University, Uli Campus, Nigeria for microbiological analyses [8-10].

### 2.3 Enrichment, Culturing and Isolation of Aromatic Hydrocarbon Degrading Bacterial Strains

The hydrocarbon degraders were isolated from sediment and water samples of the three sampling sites using modified mineral basal agar (4 g K<sub>2</sub>HPO<sub>4</sub>, 1.0 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.1 g MgSO<sub>4</sub>, 1.8 g KH<sub>2</sub>PO<sub>4</sub>, 0.1 g FeSO<sub>4</sub>, 0.1 g NaCl, 0.2 g CaCl<sub>2</sub>, 15 g Agar agar and distilled water 1,000 mL at pH 7.00 ± 0.20) enriched with xylene, anthracene and pyrene aromatic hydrocarbons. Sterilization of the medium was done by autoclaving at 121°C and 15 psi for 15 minutes. Thereafter, 0.2 mL

acetone solution containing 0.1% of the selected hydrocarbons (xylene, anthracene and pyrene) were aseptically pipetted and spread on the surface of the agar contained in Petri dish plates. The acetone was allowed to evaporate aseptically and 0.1 mL aliquots of the  $10^{-3}$  dilutions were plated on the solidified media with a glass spreader. The spreader was sterilized after each successive spreading by dipping it in 70 % ethanol and then passing it through flame of a Bunsen burner. The inoculated plates were sealed using adhesive tape and foil to prevent contamination and photolysis and later placed in black polythene bags, and then incubated in the dark at  $28.00 \pm 0.20^\circ\text{C}$  for 14 days. Also, Nutrient agar plates without hydrocarbons were inoculated and incubated at  $28.00 \pm 0.20^\circ\text{C}$  for 24 – 48 hrs [7,10-12].

## 2.4 Total Viable Count Technique

### 2.4.1 Determination of total non-hydrocarbon - utilizing bacteria (TNHUB)

The spread plate technique was used to determine the total non - hydrocarbon - utilizing bacteria on Nutrient agar. All the plates yielding 30 - 300 colonies were counted and the average

number of colonies per plates were determined. The number of total heterotrophic bacteria was expressed at CFU /g and CFU /mL [7, 12-15].

### 2.4.2 Determination of total viable hydrocarbon – utilizing bacteria (TVHUB)

Following the technique described above, total viable hydrocarbon - utilizing bacteria were determined on the modified mineral basal medium. All the plates yielding 30 - 300 colonies were counted, the average number of colonies per plates were determined and expressed at CFU /g and CFU /mL [7,12,14].

## 2.5 Purification and Maintenance of Cultures

Colonies that developed on hydrocarbon enriched plates were duplicated onto new hydrocarbon enriched plates and incubated at  $28.00 \pm 0.20^\circ\text{C}$  for 14 days. The newly developed colonies were carefully chosen as xylene, anthracene and pyrene degraders and finally sub - cultured on Bijou bottles where they are preserved at  $4^\circ\text{C}$  in refrigerator [10].




Fig. 1. Thematic map indicating the locations of sample points and its land cover in Abonema study area




Fig. 2. Thematic map indicating the locations of sample points and its land cover in Nembe study area




Fig. 3. Thematic map indicating the locations of sample points and its land cover in Onne study area

## 2.6 Screening and Selection Test

In order to screen and select the best and strongest degrading strains, the bacterial isolates were grown using 5 mL of all the isolates in test tubes comprising 25 mL of the modified mineral basal medium supplemented with 100 ppm of xylene, anthracene and pyrene hydrocarbons as described in sub-section 2.3 above. Thereafter, the set ups were incubated at  $28.00 \pm 2.00^\circ\text{C}$  for five days. Bacteria that started growing fast with high turbidity in the vicinity of the medium containing aromatic compounds, quantified at 600 nm using a UV - VIS spectrophotometer (Astell, UV - Vis Grating, 752 W) were selected as xylene, anthracene and pyrene degraders [10, 16,17].

## 2.7 Characterization and Identification of Selected Hydrocarbon Utilizing Bacterial Isolates

### 2.7.1 Molecular characteristics

#### 2.7.1.1 Identification of bacteria

The Gram - reaction test was first used to verify the morphological characterization of the bacterial colonies prior to molecular identification [18].

#### 2.7.1.2 Genomic DNA extraction of the bacterial isolates

DNA extraction was conducted using conventional method of Cetyltrimethyl Ammonium Bromide (CTAB) protocol in sterile Eppendorf tubes. The DNA was separated electrophoretically with 1 % agarose gel containing 0.1  $\mu\text{g}$  /mL of ethidium bromide stain. The 20 cm long gels were ran at 80 V /cm for 60 minutes in 100 mL of 1x tris acetate EDTA (TAE) electrophoresis buffer. The DNA was visualised by UV fluorescence to determine the success of the extraction process [18].

#### 2.7.1.3 Polymerase chain reaction (PCR) and sequencing of the extracted DNA

The master mix aliquot for the PCR was dispensed into individual PCR tube and the different DNA samples were added to each tubes. The negative control was used to check for contamination in the master mix. The PCR reagents in each tube amounted to 50  $\mu\text{L}$  containing: buffer (5  $\mu\text{L}$ ),  $\text{MgCl}_2$  (1.5  $\mu\text{L}$ ), primer 1 (forward 16S - P1 PCR 5' -

AGAGTTTGATCCTGGCTCAG - 3') (2  $\mu\text{L}$ ), primer 2 (reverse 16S - P2 PCR 5' - AAGGAGGTGATCCAGCCGCA - 3') (2  $\mu\text{L}$ ), dNTP mix (1  $\mu\text{L}$ ), Dream Taq (0.25  $\mu\text{L}$ ), sterile sabax water (35.25  $\mu\text{L}$ ) and purified DNA samples (3  $\mu\text{L}$ ). The PCR reactions was performed using MJ Mini thermal cycler (Bio-Rad, Hercules, CA, USA). The cycling conditions were set at (a) initial denaturation 10 minutes at  $95^\circ\text{C}$  for 1 cycle. (b) Denaturation at  $95^\circ\text{C}$  for 30 seconds, (c) Annealing cycling at  $94^\circ\text{C}$  for 30 seconds, (d) Elongation at  $54^\circ\text{C}$  for 2 minutes. All steps in denaturation, annealing and elongation was for 35 cycles and (e) final elongation 10 minutes at  $72^\circ\text{C}$  for 1 cycle. The reaction was held at  $4^\circ\text{C}$  for 1 hr in the thermal cycler. The The PCR products (genes) were separated at 1 kbp size and visualized as stated above (Fig. 1) [18]. Then the PCR products (20  $\mu\text{L}$  each) were cleaned up later using 160  $\mu\text{L}$  of 13% polyethylene glycol (PEG) 8000, 20  $\mu\text{L}$  of 5 M NaCl solution and 200  $\mu\text{L}$  of 70% ethanol. The cleaned PCR products were sent for sequencing and was conducted using the automated DNA sequencer (Perkin-Elmer) in line with the sequencing kit protocol at the Forestry and Agricultural Biotechnology Institute (FABI) Sequencing Facility, University of Pretoria, South Africa [18].

#### 2.7.1.4 Blasting and phylogenetic correctional analyses

The blasting of DNA sequences was performed by revising the sequences of the 16S rRNA genes obtained using BioEdit software, corrected sequences duplicated in a FASTA format form and completed in the National Centre for Biotechnology Information (NCBI) website. Homologies of the gene sequences were checked and compared with the sequences of the NCBI database and finally aligned using MAFFT software [19]. The taxonomic correctional studies were done using Mega 7 software and evolutionary distance of the isolates were calculated using neighbour - joining (NJ) methods [4,15,18,19].

#### 2.7.1.5 GenBank accession number

NCBI accession numbers were assigned to the nine selected aromatic hydrocarbon degrading bacterial strains and the nucleotide sequence details of 16S rRNA genes of these strains have been submitted to the NCBI /GenBank database since 01/12/16 under the accession numbers from KY171979 - KY171987 [3,4,15,19-21].

**Table 1. Mean total non – hydrocarbon utilizing bacterial (TNHUB) count of sediment and water samples from the three sampled locations**

Samples	Log CFU /g/ml	Percentage (%)
Abonema water	13.60 ± 0.02	20.80 ± 0.16
Abonema sediment	12.70 ± 0.02	18.90 ± 0.20
Nembe water	18.95 ± 0.04	32.40 ± 0.16
Nembe sediment	11.50 ± 0.04	16.30 ± 0.16
Onne water	08.00 ± 0.16	08.70 ± 0.20
Onne sediment	05.34 ± 0.02	02.90 ± 0.06

*The values are mean ± standard deviation of triplicate determination*

**Table 2. Mean total viable hydrocarbon utilizing bacterial (TVHUB) count of sediment and water samples from the three sampled locations**

Sample	Log CFU /g/ml	Percentage (%)
Abonema water + xylene	04.35 ± 0.04	02.50 ± 0.04
Abonema sediment + xylene	30.20 ± 0.12	17.30 ± 0.12
Abonema water + anthracene	09.20 ± 0.08	05.30 ± 0.08
Abonema sediment + anthracene	08.05 ± 0.01	04.60 ± 0.02
Abonema water + pyrene	04.42 ± 0.04	02.50 ± 0.04
Abonema sediment + pyrene	09.45 ± 0.01	05.40 ± 0.02
Nembe water + xylene	12.35 ± 0.02	07.10 ± 0.06
Nembe sediment + xylene	09.50 ± 0.01	05.50 ± 0.01
Nembe water + anthracene	10.50 ± 0.01	06.00 ± 0.12
Nembe sediment + anthracene	06.09 ± 0.03	03.50 ± 0.02
Nembe water + pyrene	15.35 ± 0.15	08.80 ± 0.02
Nembe sediment + pyrene	08.25 ± 0.02	04.70 ± 0.02
Onne water + xylene	07.60 ± 0.01	04.40 ± 0.02
Onne sediment + xylene	07.70 ± 0.08	04.40 ± 0.08
Onne water + anthracene	05.20 ± 0.02	03.00 ± 0.06
Onne sediments + anthracene	08.50 ± 0.02	04.90 ± 0.01
Onne water + pyrene	10.50 ± 0.02	06.00 ± 0.02
Onne sediment + pyrene	06.95 ± 0.12	04.00 ± 0.12

*The values are mean ± standard deviation of triplicate determination*

**Table 3. Growth performance of the aromatic hydrocarbon - degraders isolated from Abonema sampled location**

Isolate	Optical density (OD <sub>600 nm</sub> )		Pyrene
	Xylene	Anthracene	
AB1	0.657 ± 0.008	0.657 ± 0.001	0.580 ± 0.003
ANT4*	0.676 ± 0.004	0.775 ± 0.007	0.822 ± 0.002
AB3	0.701 ± 0.021	0.467 ± 0.029	0.666 ± 0.003
AB4	0.715 ± 0.004	0.598 ± 0.024	0.841 ± 0.001
PYR3*	0.598 ± 0.005	0.511 ± 0.003	1.041 ± 0.008
AB6	0.641 ± 0.001	0.494 ± 0.002	0.653 ± 0.001
AB7	0.687 ± 0.004	0.690 ± 0.002	0.803 ± 0.016
AB8	0.618 ± 0.001	0.638 ± 0.007	0.782 ± 0.003
AB9	0.457 ± 0.002	0.475 ± 0.001	0.573 ± 0.004
XYL2*	0.952 ± 0.004	0.312 ± 0.002	0.838 ± 0.021
AB11	0.793 ± 0.014	0.495 ± 0.002	0.970 ± 0.003
AB12	0.647 ± 0.002	0.446 ± 0.001	0.621 ± 0.005
AB13	0.328 ± 0.001	0.415 ± 0.001	0.451 ± 0.001

\* = Isolates with highest degradability; values are mean ± standard deviation of triplicate determination

## 2.8 Detection of Catabolic and Surfactant Genes by PCR Analyses

### 2.8.1 PCR primer design

Following the method of Swaathy et al. [19] as modified in this study, a portion of catechol gene 881 bp (C23O) was pulled out from the genomic DNA using F: 5'- ATG AGC AAC AAA TAC GAA TT- 3' and R: 5'- TCA AAC GGT CAA TCT GAT AT- 3' (Fig. 6). Likewise, according to the method of Qazi et al. [21] and Swaathy et al. [3] as modified in this study, the primer pair of microsurf gene *srfA3* /*licA3* using F: CAAAACGCAKCATATGAG and *srfA3* /*licA3* and R: AGCGGAYATATTGATGCGGYTC was designed to amplify a 881 bp portion of the *srfA3* or the homologous *licA3* gene present in surfactin/lichenysin and gene - specific primers of rhamnolipid gene (*rhlB*) using kpd - F 5'- GCCACGACCAGTTCGAC-3' and kpd - R 5' CATCCCCCTCCCTATGAC-3' (Fig. 6). They were subjected to the same PCR and gel electrophoretic conditions and protocols stated above [18].

### 2.9 Data Analysis

All values were expressed as mean  $\pm$  S.D. and ordinary one-way analysis of variance (ANOVA) followed by post Tukey's, multiple comparison test was performed on the data obtained using Graph-Pad Prism version 7.00. The results were considered statistically significant at 95% confidence intervals ( $P = .05$ ) [15,16].

## 3. RESULTS

### 3.1 Bacterial Enumeration

The result of the mean total non-hydrocarbon utilizing bacterial (TNHUB) count of sediment and water samples from the three sampled locations is presented on Table 1. From the result, Nembe water had the highest significant ( $P = .05$ ) mean TNHUB count of log 18.95  $\pm$  0.04 CFU /mL with highest percentage of occurrence (32.40  $\pm$  0.16 %) while Onne sediment had the lowest significant ( $P = .05$ ) mean TNHUB count of log 05.34  $\pm$  0.02 CFU /g with lowest percentage of occurrence (02.90  $\pm$  0.06 %) respectively. Similarly, the result of the mean total viable hydrocarbon utilizing bacterial (TVHUB) count sediment and water samples from the three sampled locations is presented on Table 2. From the result, Abonema sediment with xylene and

Abonema water with xylene had the highest significant ( $P = .05$ ) mean TVHUB count of log 30.20  $\pm$  0.12 CFU /g with highest percentage of occurrence (17.30  $\pm$  0.12%) and the lowest significant ( $P = .05$ ) mean TVHUB count of log 04.35  $\pm$  0.04 CFU /mL with lowest percentage of occurrence (02.50  $\pm$  0.04%) respectively.

### 3.2 Degradability Test

The result of the growth performance (OD<sub>600</sub> nm) of the aromatic hydrocarbon - degraders isolated from Abonema sampled location is presented in Table 3. From the result, 13 isolates were obtained with strains XYL2, ANT4 and PYR3 having the highest significant ( $P = .05$ ) absorbance values of 0.952  $\pm$  0.004, 0.775  $\pm$  0.007 and 1.041  $\pm$  0.008 on xylene, anthracene and pyrene hydrocarbons. The result of the growth performance (OD<sub>600</sub> nm) of the aromatic hydrocarbon - degraders isolated from Nembe sampled location is presented in Table 4. From the result, 17 isolates were obtained with strains XYL7, ANT1 and PYR5 having the highest significant ( $P = .05$ ) absorbance values of 1.055  $\pm$  0.002, 0.816  $\pm$  0.007 and 0.933  $\pm$  0.007 on xylene, anthracene and pyrene hydrocarbons. The result of the growth performance (OD<sub>600</sub> nm) of the aromatic hydrocarbon - degraders isolated from Onne sampled location is presented in Table 5. From the result, 18 isolates were obtained with strains XYL8, ANT6 and PYR9 having the highest significant ( $P = .05$ ) absorbance values of 0.741  $\pm$  0.007, 1.433  $\pm$  0.013 and 0.871  $\pm$  0.001 on xylene, anthracene and pyrene hydrocarbons. On the basis of these results, strains ANT1, XYL2, PYR3, ANT4, PYR5, ANT6, XYL7, XYL8 and PYR9 were selected as the best and strongest degraders of xylene, anthracene and pyrene hydrocarbons.

### 3.3 Molecular Characterization of Bacterial Isolates

The result of the PCR - amplification of 16S rRNA genes of the aromatic degrading bacteria genomic DNA is shown in Fig. 4. From the result, it was confirmed that the PCR products obtained using universal primer 16S - P1 PCR (27 F 5' - : AGA GTT TGA TCC TGG CTC AG - 3') and 16S - P2 PCR (1492 R 5' - ACG GCT ACC TTG TTA CGA CTT - 3') have the molecular weight of 1 kbp visualized by UV fluorescence under agarose gel electrophoresis. The result of the percentage similarity and GenBank accession numbers of 16S rRNA sequences of the closest relative for the aromatic degrading bacterial

isolates is presented in Table 6. From the result, *Alcaligenes faecalis* was the most blasted bacteria with high similarity (98 - 99%) followed by *Providencia* spp. (95 - 97%), *Brevundimonas diminuta* (100%), *Myroides odoratus* (90%), *Serratia marcescens* (97%) and *Bacillus cereus* (98%) using NCBI BLAST software. The result of neighbour - joining phylogenetic relationship among the 16S rRNA sequence of the aromatic degrading bacterial isolates constructed by MEGA 7.0 is shown in Fig. 5. From the result, it revealed that the nine bacterial isolates (KY171979, KY171984, KY171987, KY171980, KY171982, KY171981, KY171985, KY171986 and KY171983) arise from similar clusters and node establishing that they are correlationally connected to one another with common origin.

### 3.4 Detection of Catabolic and Surfactant Genes

The result of the catabolic and surfactant genes detected in some of the marine aromatic degrading bacterial isolates using specific primers for each gene is presented in Table 7 while the result of the PCR – amplification of primers specific for catabolic gene (C23O) and surfactant genes (*rhIB*, *SrfA3* /*LicA3*) of *Providencia vermicola* strain ANT1, *Alcaligenes faecalis* strain XYL2, *Serratia marcescens* strain XYL7 and *Providencia* sp. strain XYL8 is shown in Fig. 6. From the results, catabolic gene

(C23O) was detected in only four (4) out of the nine marine aromatic degrading bacteria with name stated above while surfactant genes (*rhIB*, *SrfA3* /*LicA3*) were detected only in three (3) out of the nine marine aromatic degrading bacteria all of which had 881 base pairs sizes of PCR products of the catabolic and surfactant genes visualized by UV fluorescence under agarose gel electrophoresis respectively.

## 4. DISCUSSION

In this study, an attempt was made to explore the diversity, catabolic and surfactant genes of marine hydrocarbon utilizing bacteria from our three study areas and the result in Tables 1 and 2 showed that surface water harbours more non – hydrocarbon utilizing bacteria which could possibly be due to the more nutrient and oxygen levels in surface water than the sediment samples. The result in Table 2 showed that sediment surface harbours more hydrocarbon utilizing bacteria than surface water possibly due to the high accumulative nature of sediment to aromatic compounds than water. Aromatic compounds tend to have low solubility to water hence making them adsorbed more to sediment than surface water. Comparatively, there was significant abundance of NHUB more than HUB and more xylene degraders than anthracene and pyrene degraders in the both samples respectively. The result uphold the findings of

**Table 4. Growth performance of the aromatic hydrocarbon - degraders isolated from Nembe sampled location**

Isolate	Optical density (OD <sub>600</sub> nm)		Pyrene
	Xylene	Anthracene	
NW1	0.885 ± 0.003	0.236 ± 0.005	0.708 ± 0.008
PYR5*	0.710 ± 0.003	0.216 ± 0.005	0.933 ± 0.007
NW3	0.466 ± 0.007	0.201 ± 0.000	0.806 ± 0.004
NW4	0.893 ± 0.002	0.356 ± 0.008	0.827 ± 0.008
NW5	0.750 ± 0.004	0.132 ± 0.005	0.767 ± 0.008
NW6	0.644 ± 0.004	0.246 ± 0.004	0.724 ± 0.008
NW7	0.561 ± 0.003	0.193 ± 0.005	0.808 ± 0.001
NW8	0.628 ± 0.008	0.472 ± 0.001	0.826 ± 0.008
XYL7*	1.055 ± 0.002	0.588 ± 0.005	0.927 ± 0.001
NW10	0.809 ± 0.002	0.785 ± 0.002	0.881 ± 0.004
NW11	0.826 ± 0.001	0.444 ± 0.002	0.891 ± 0.001
NW12	0.625 ± 0.005	0.563 ± 0.001	0.728 ± 0.006
NW13	0.374 ± 0.008	0.775 ± 0.001	0.760 ± 0.001
NW14	0.701 ± 0.001	0.622 ± 0.003	0.788 ± 0.007
NW15	0.705 ± 0.008	0.529 ± 0.004	0.830 ± 0.002
NW16	0.769 ± 0.002	0.380 ± 0.001	0.822 ± 0.001
ANT1*	0.804 ± 0.003	0.816 ± 0.007	0.583 ± 0.001

\* = Isolates with highest degradability; values are mean ± standard deviation of triplicate determination

**Table 5. Growth performance of the aromatic hydrocarbon - degraders isolated from Onne sampled location**

Isolate	Optical density (OD <sub>600</sub> nm)		Pyrene
	Xylene	Anthracene	
ON1	0.721 ± 0.001	0.884 ± 0.007	0.500 ± 0.001
ON2	0.204 ± 0.001	0.660 ± 0.011	0.454 ± 0.001
ON3	0.473 ± 0.003	0.476 ± 0.036	0.561 ± 0.013
ON4	0.207 ± 0.001	0.766 ± 0.001	0.565 ± 0.033
ON5	0.477 ± 0.002	0.457 ± 0.001	0.378 ± 0.005
ON6	0.409 ± 0.005	0.489 ± 0.100	0.562 ± 0.021
ON7	0.251 ± 0.003	0.428 ± 0.014	0.728 ± 0.001
ON8	0.111 ± 0.005	0.429 ± 0.014	0.425 ± 0.021
ON9	0.463 ± 0.008	0.357 ± 0.011	0.281 ± 0.006
PYR9*	0.106 ± 0.001	0.335 ± 0.001	0.871 ± 0.001
ON11	0.700 ± 0.001	0.901 ± 0.005	0.417 ± 0.002
ANT6*	0.511 ± 0.006	1.433 ± 0.013	0.568 ± 0.009
ON13	0.273 ± 0.002	0.386 ± 0.002	0.527 ± 0.001
ON14	0.278 ± 0.005	0.553 ± 0.022	0.684 ± 0.003
ON15	0.291 ± 0.003	0.748 ± 0.009	0.522 ± 0.010
ON16	0.662 ± 0.001	0.919 ± 0.002	0.494 ± 0.002
XYL8*	0.741 ± 0.007	0.510 ± 0.013	0.602 ± 0.004
ON18	0.354 ± 0.002	1.004 ± 0.001	0.478 ± 0.001

\* = Isolates with highest degradability; values are mean ± standard deviation of triplicate determination

**Table 6. Percentage similarity and GenBank accession numbers of 16S rRNA sequences of the closest relative for the aromatic degrading bacterial isolates**


Isolate code	Closest relative in GenBank	Max score	Total score	Query coverage	E. value	Max identity	Accession number
ANT1	<i>Providencia vermicola</i> strain MTCC 5578	544	544	93%	6e-151	95%	KY171979
XYL2	<i>Alcaligenes faecalis</i> strain MOR02	1559	4679	99%	0.0	99%	KY171984
PYR3	<i>Brevundimonas diminuta</i> strain zjs01	1489	1489	97%	0.0	100%	KY171987
ANT4	<i>Alcaligenes faecalis</i> strain MOR02	1594	1594	99%	0.0	99%	KY171980
PYR5	<i>Alcaligenes faecalis</i> strain MOR02	1537	4612	98%	0.0	98%	KY171982
ANT6	<i>Myroides odoratus</i> strain D25T	1194	1194	95%	0.0	90%	KY171981
XYL7	<i>Serratia marcescens</i> strain SM6	1476	1476	98%	0.0	97%	KY171985
XYL8	<i>Providencia</i> sp. strain X1	1491	1491	98%	0.0	97%	KY171986
PYR9	<i>Bacillus cereus</i> strain B4	1543	16940	98%	0.0	98%	KY171983

Chikere et al. [7] who reported that these waterways are unceasingly open to petroleum hydrocarbons due to navigational actions and this possibly will have augmented the sediment with hydrocarbon utilizing bacteria. However, the insufficiency of the HUB counts could be ascribed to the lack of nutrients at that depth especially nitrogen and phosphorus which reduce with input of hydrocarbons. Additional factor that decreases existing metabolic nutrients in marine ecosystem according to Xu et al. [22], is substantial percolation initiated by tidal flood and wave action. The existence of hydrocarbon utilizers in the midst of the heterotrophic population in the samples is a sign of earlier contamination owing to hydrocarbon pollution [12,23].

**Table 7. Catabolic and surfactant genes detected in some of the marine aromatic degrading bacterial isolates using specific primers for each gene**

Isolate code	Bacterial name	Expected band (bp)	C23O	rhIB	SrfA3 /LicA3
ANT1	<i>Providencia vermicola</i>	881 bp	+	-	-
XYL2	<i>Alcaligenes faecalis</i>	881 bp	+	+	-
PYR3	<i>Brevundimonas diminuta</i>	-	-	-	-
ANT4	<i>Alcaligenes faecalis</i>	-	-	-	-
PYR5	<i>Alcaligenes faecalis</i>	-	-	-	-
ANT6	<i>Myroides odoratus</i>	-	-	-	-
XYL7	<i>Serratia marcescens</i>	881 bp	+	+	+
XYL8	<i>Providencia sp.</i>	881 bp	+	+	+
PYR9	<i>Bacillus cereus</i>	-	-	-	-

+ Sign indicates the PCR product was detected and – sign means PCR product was not detected




**Fig. 4. PCR - amplification of 16S rRNA genes of the aromatic degrading bacteria genomic DNA**

Following isolation, a total of nine (9) isolates ANT1, XYL2, PYR3, ANT4, PYR5, ANT6, XYL7, XYL8 and PYR9 out of the 48 isolates (9/48) representing 18.75 % of the isolates were screened and selected as best and strongest degraders of xylene, anthracene and pyrene hydrocarbons which they significantly ( $P = .05$ ) utilize as source of carbon and energy and are indicated by absorbance values of each isolates (Tables 3, 4 and 5). The result is justified by the finding of Mao et al. [24], who reported that the enrichment and isolation of native PAH degraders is the most essential feature of microbial degradation of PAH as the native PAH degraders possess the adaptive traits to these hydrocarbons. Pathak and Bhatnagar [25],

reported that enrichment culturing is very imperative towards the attainment of hydrocarbon bioremediation as it leads to selection of microbial degraders that are better adapted to hydrocarbons. Esedafe et al. [12] reported that an occurrence of 3/41 representing 7.32% isolates from refinery effluent were efficient in degrading the polyaromatic hydrocarbons (PAHs).

The result in Fig. 4 confirmed that the PCR products of the genomic DNA of the isolated bacteria had the molecular weight of 1 kbp. Akinbankole et al. [26], obtained 1,500 bp PCR product from anthracene and pyrene degrading bacteria isolated from petroleum polluted water




**Fig. 5. Neighbour - joining phylogenetic relationship among the 16S rRNA sequence of the aromatic degrading bacterial isolates constructed by MEGA 7.0**

Numbers at the nodes indicate bootstrap support (%) based on 1000 replicates. The sum of branch length = 0.93646865 using p - distance method involving 9 nucleotide sequences with total of 303 positions. GenBank accession numbers are given in parentheses

and soil in Malaysia in their research finding. Yuliani et al. [17], obtained in their research 1,489 bp PCR product from phenanthrene and pyrene degrading bacteria isolated from marine region of Indonesia. Isiodu et al. [15], reported that all the seven (7) polyaromatic hydrocarbon

utilizing bacterial isolates isolated from Bodo Creek brackish water in Nigeria showed amplification with an amplicon size of 500 bp. The result in Table 6 showed that *Alcaligenes faecalis* was the most occurring organism after blasting with high sequence similarity (98 - 99%)



**Fig. 6. PCR – amplification of primers specific for catabolic gene (C230) and surfactant genes (rhIB, SrfA3 /LicA3) of *Providencia vermicola* strain ANT1, *Alcaligenes faecalis* strain XYL2, *Serratia marcescens* strain XYL7 and *Providencia* sp. strain XYL8**

followed by *Providencia* spp. (95 - 97%), *Brevundimonas diminuta* (100%), *Myroides odoratus* (90%), *Serratia marcescens* (97%) and *Bacillus cereus* (98%) using NCBI BLAST software. Akinbankole et al. [26], found out that *B. cereus* was the most blasted organism with 99% sequence similarity. The phylogenetic tree showed the evolutionary correlation among hydrocarbon metabolizing bacteria isolated in this study. The result shown in Fig. 5 revealed that the nine bacterial isolates (KY171979, KY171984, KY171987, KY171980, KY171982, KY171981, KY171985, KY171986 and KY171983) had similar lineage as they arise from analogous node and hence they are evolutionary connected to their relatives in the GenBank. The sequences of these genes have been submitted to NCBI/GenBank database since first of December two thousand and sixteen (01/12/16). They belong to the genera: *Providencia*, *Alcaligenes*, *Brevundimonas*, *Myroides*, *Serratia*, and *Bacillus*; and families of: Enterobacteriaceae, Alcaliginaceae, Caulobacteriaceae, Flavobacteriaceae, and Bacillaceae; and phyla of: Proteobacteria, Bacteroidetes and Firmicutes which members have been implicated in petroleum and aromatic hydrocarbon biodegradation by several authors [7,10,15,17,19,26-33].

In order to broaden the description of the aromatic hydrocarbon - degrading bacteria isolated from the Rivers marine environment, PCR assays were performed to explore for

functional genes coding for the catechol 2, 3 - dioxygenase enzyme (C230) and genes related to surfactant production namely rhamnolipid enzyme (*rhIB*) and surfactin/lichenysin enzyme (*SrfA3 /LicA3*) and the results are presented and shown in Table 7 and Fig. 6. From the results, catabolic gene (C230) was detected in only four (4) out of the nine marine aromatic degrading bacteria with name stated above while surfactant genes (*rhIB*, *SrfA3 /LicA3*) were detected only in three (3) out of the nine marine aromatic degrading bacteria all of which had 881 base pairs sizes of PCR products of the catabolic and surfactant genes. Even though biosurfactants aid in solubilizing or mediating the interface between the bacterial degraders and the aromatic compounds, the catabolic responses detected in our study were effected by the dioxygenase genes as revealed from the amplified products of 881 bps. The detection of C230 gene in some of the Gram negative bacteria and not in the Gram positive bacteria showed that these strains (ANT1, XYL2, XYL7 and XYL8) synthesize dioxygenases for either complete or partial breakdown of aromatic hydrocarbons (xylene, anthracene and pyrene) and a probable reason could be that the detected genes were highly preserved among diverse Gram - negative bacteria; hence support the research work of Hesham et al. [4], who confirmed the existence of both monooxygenase and dioxygenase in *S. koreensis* strain ASU - 06. C12O and C23O dioxygenases have been known to contribute a significant part in the catabolism of aromatic

rings by the bacteria as they are responsible for splitting of aromatic C – C bond at ortho or meta positions. The implication of the gene C23O as an essential gene in the catabolism of low and high molecular weights PAHs has been reported by Swaathy et al. [19]. Moreso, the detection of rhamnolipid (*rhIB*) and surfactin /lichenysin (*SrfA3 /LicA3*) genes in these strains support the continuous argument among the scientists about the occurrence of *rhIB* and *SrfA3 /LicA3* genes in bacteria other than *Pseudomonas* and *Bacillus* species and several new strains of bacteria having exceptional ability to synthesize rhamnolipids have in recent times been described. There are dearth of reports documented on rhamnolipid production among pathogens especially *Burkholderia mallei* and *B. pseudomallei* and the non - pathogenic especially *B. thailandensis*. This observation contradicts the findings of Swaathy et al. [3,19], who found out that the expression of *srf* and *licA3* genes were only realized in all of the five *Bacillus* species and marine *Bacillus licheniformis* MTCC 5514. On the other hand, those isolates that could grow on the aromatic hydrocarbons but had unsuccessful amplification, could be as a result of incompatibilities among the tested primers and gene sequences [34] and may contain other catabolic and surfactant genes aside the tested genes.

## 5. CONCLUSION

The whole study revealed that the three sampling sites harbour a lot of phylogenetically related aromatic hydrocarbon degrading bacterial strains belonging to the genera: *Providencia*, *Alcaligenes*, *Brevundimonas*, *Myroides*, *Serratia*, and *Bacillus* which are able to utilize xylene, anthracene and pyrene as source of carbon and energy. It also revealed that some of the isolated bacteria especially *Serratia marcescens* XYL7 possess functional genes coding for aromatic degradation and surfactant production. Our results suggest that these bacteria may possibly play a significant part in degradation of simple, low and high molecular weight aromatic hydrocarbons and could be recommended for *in situ* aromatic hydrocarbon bioremediation in the Niger Delta environment.

## ACKNOWLEDGEMENTS

The research work described in this paper was supported by fund from the year 2012/2013 (Merged) Tertiary Education Trust Fund (TETFUND) Academic Staff Training and

Development of Chukwuemeka Odumegwu Ojukwu Universit, Uli Campus. Also, I want to thank in a special way Prof. (Dame) E. I. Chukwura who supervise the work and Mr Ubani, Onyedika of Department of Environmental Sciences, University of South Africa, Pretoria, South Africa for his technical assistance towards completion of the work. The anonymous reviewers are sincerely thanked for their beneficial suggestions to improve the manuscript.

## COMPETING INTERESTS

Author has declared that no competing interests exist.

## REFERENCES

1. Nwilo PC, Badejo OT. Oil spill problems and the management in the Niger Delta. International Oil Spill Conference, Miami, Florida, USA. 2005;20-2.
2. Rajaei S, Seyedi SM, Raiesi F, Shiran B, Raheb J. Characterization and potentials of oil-degrading bacteria inhabiting the rhizosphere of wild oat (*Avena fatua* L.) in South West of Iran. Iran J Biotech. 2013; 11(1):32- 40.
3. Swaathy S, Kavitha V, Pravin AS, Sekaran G, Mandal AB, Gnanamani A. (2014a). Phylogenetic framework and biosurfactant gene expression analysis of marine *Bacillus* spp. of Eastern Coastal Plain of Tamil Nadu. Internat J Bact. 2014a; 860491.
4. Hesham AE, Mawad AMM, Mostafa YM, Shoreit A. Biodegradation ability and catabolic genes of petroleum-degrading *Sphingomonas koreensis* strain ASU-06 isolated from Egyptian oily soil. BioMed Res Internat. 2014;127674.
5. Nakamura FM, Germano MG, Tsai SM. Capacity of aromatic compound degradation by bacteria from Amazon Dark Earth. Diver. 2014;6:339-53.
6. Ma J, Xu L, Jia L. Degradation of polycyclic aromatic hydrocarbons by *Pseudomonas* sp. JM2 isolated from active sewage sludge of chemical plant. J Environ Sci. 2012;24(12):2141-8.
7. Chikere CB, Okpokwasili GC, Ichiakor O. Characterization of hydrocarbon utilizing bacteria in tropical marine sediments. Afr J Biotech. 2009;8(11):2541-44.
8. Abu GO, Chikere BO. Cell surface properties of hydrocarbon utilizing bacterial

- isolates from the Port Harcourt marine environment. *Nig J Microbiol.* 2006;20(1): 809-16.
9. Gorleku MA, Carboo D, Palm LMN, Quasie WJ, Armah AK. Polycyclic aromatic hydrocarbons (PAHs) pollution in marine waters and sediments at the Tema Harbour, Ghana. *Acad J Environ Sci.* 2014;2(7):108-15.
  10. John RC, Essien JP, Akpan SB, Okpokwasili GC. Polycyclic aromatic hydrocarbon degrading bacteria from aviation fuel spill site at Ibeno, Nigeria. *Bull Environ Contam Toxicol.* 2012;88:1014-9.
  11. John RC, Okpokwasili GC. Crude oil-degradation and plasmid profile of nitrifying bacteria isolated from oil impacted mangrove sediment in the Niger Delta of Nigeria. *Bull Environ Contam Toxicol.* 2012;88:1020-6.
  12. Esedafe WK, Fagade OE, Umaru FF, Akinwotu O. Bacterial degradation of the polycyclic aromatic hydrocarbon (PAH) - fraction of refinery effluent. *Internat J Environ Biorem Biodeg.* 2015;3(1):23-7.
  13. Bahig AE, Aly EA, Khaled AA, and Amel KA. Isolation, characterization and application of bacterial population from agricultural soil at Sohag Province, Egypt. *Mal J Microbiol.* 2008;4(2):42-50.
  14. Kafilzadeh F, Pour FH, Tahery Y, Azad HN. Bioremediation of pyrene by isolated bacterial strains from the soil of the landfills in Shiraz (Iran). *Ann Biol Res.* 2012;3(1): 486-94.
  15. Isiodu GG, Stanley HO, Ezebuio V, Okerentugba PO. Role of plasmid-borne genes in the biodegradation of polycyclic aromatic hydrocarbons (PAHs) by consortium of aerobic heterotrophic bacteria. *J Petr Environ Biotech.* 2016;7(1): 264.
  16. Kafilzadeh F, Pour FH. Degradation of naphthalene, phenanthrene and pyrene by *Pseudomonas* sp. and *Corynebacterium* sp. in the landfills. *Internat J Biosci.* 2012; 2(9):77-84.
  17. Yuliani H, Sahlan M, Hermansyah H, Wijanarko A. Selection and identification of polyaromatic hydrocarbon degrading bacteria. *World Appl Sci J.* 2012;20(8): 1133-8.
  18. Ubani O, Atagana HI, Thantsha MS, Rasheed A. Identification and characterisation of oil sludge degrading bacteria isolated from compost. *Arch Environ Prot.* 2016;42(2):67-77.
  19. Swaathy S, Kavitha V, Pravin AS, Mandal, AB, Gnanamani A. Microbial surfactant mediated degradation of anthracene in aqueous phase by marine *Bacillus licheniformis* MTCC 5514. *Biotech Rep.* 2014b;4:161-70.
  20. Guo C, Dang Z, Wong Y, Tam NF. Biodegradation ability and dioxygenase genes of PAH-degrading *Sphingomonas* and *Mycobacterium* strains isolated from mangrove sediments. *Internat Biodet Biodeg.* 2010;64:419-26.
  21. Qazi MA, Malik ZA, Qureshi GD, Hameed A, Ahmed S. Yeast extract as the most preferable substrate for optimized biosurfactant production by *rhlB* gene positive *Pseudomonas putida* SOL-10 isolate. *J Biorem Biodeg.* 2013;4(7):204.
  22. Xu R, Lau NLA, Ng KL, Obbard JP. Application of a slow-release fertilizer for oil bioremediation in beach sediments. *J Environ Qual.* 2004;33:1210-6.
  23. Al-Thani RF, Abd-El-Haleem DAM, Al-Shammri M. Isolation and characterization of polyaromatic hydrocarbons-degrading bacteria from different Qatari soils. *Afr. J. Microbiol. Res.* 2009;3(11):761-6.
  24. Mao J, Luo Y, Teng Y, Li Z. Bioremediation of polycyclic aromatic hydrocarbon-contaminated soil by a bacterial consortium and associated microbial community changes. *Internat Biodet Biodeg.* 2012;70:141-7.
  25. Pathak H, Bhatnagar K. *Alcaligenes*-the 4T engine oil degrader. *J Biorem and Biodeg.* 2011;2(4):2-5.
  26. Akinbankole AS, Tunung R, Tennant AM. Biochemical and molecular characterization of pyrene and anthracene metabolizing bacteria isolated from oil contaminated water and soil in Malaysia. *J Appl Environ Microbiol.* 2015;3(1):25-30.
  27. Amer RA, Mapelli F, El-Gendi HM, Barbato M, Goda DA, Corsini A, Cavalca L, Fusi M, Borin S, Daffonchio D, Abdel-Fattah YR. Bacterial diversity and bioremediation potential of the highly contaminated marine sediments at El-Max District (Egypt, Mediterranean Sea). *Biomed Res Internat* 2015;2015:981-9.
  28. Fagbemi OK, Sanusi AI. Chromosomal and plasmid mediated degradation of crude oil by *Bacillus coagulans*, *Citrobacter koseri* and *Serratia ficaria* isolated from the soil. *Afr J Biotech.* 2017;16(21):1242-53.
  29. Athar MA, Akbar A, Khan YH, Ali I, Mehmood U, Sabri AN, Hasnain S.

- Characterization of hydrocarbon degrading bacteria isolated from hydrocarbon contaminated soil. J Pure Appl Microbiol. 2014;8(6):1-9.
30. Ichor T, Okerentugba PO, Okpokwasili GC. Biodegradation of total petroleum hydrocarbon by aerobic heterotrophic bacteria isolated from crude oil contaminated brackish waters of Bodo Creek. J Biorem Biodeg. 2014;5(5):236-41.
31. Irshaid FI, Jacob JH. Screening and characterization of aerobic xylene-degrading bacteria from gasoline contaminated soil sites around gas stations in Northern Jordan. J Biol Sci. 2015;15(4): 167-76.
32. Louvado A, Gomes NCM, Simões MMQ, Almeida A, Cleary DFR, Cunha A. Polycyclic aromatic hydrocarbons in deep sea sediments: Microbe–pollutant interactions in a remote environment. Sci Tot Environ. 2015;526:312-28.
33. Patowary K, Saikia RR, Kalita MC, Deka S. Degradation of polyaromatic hydrocarbons employing biosurfactant-producing *Bacillus pumilus* KSs2. Ann Microbiol. 2015;65: 225-34.
34. Ding GC, Heuer H, Zühlke S, Spiteller M, Pronk GJ, Heister K et al. Soil type-dependent responses to phenanthrene as revealed by determining the diversity and abundance of polycyclic aromatic hydrocarbon ring - hydroxylating dioxygenase genes by using a novel PCR detection system. Appl Environ Microbiol. 2010;76(14):4765-71.

© 2018 Uba; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/46533>